

THE ARENS REGULARITY AND WEAK TOPOLOGICAL CENTER OF MODULE ACTIONS

KAZEM HAGHNEJAD AZAR

ABSTRACT. Let A be a Banach algebra and A^{**} be the second dual of it. We define $\tilde{Z}_1(A^{**})$ as a weak topological center of A^{**} with respect to first Arens product and find some relations between it and the topological center of A^{**} . We also extend this new definition into module actions and find relationship between weak topological center of module actions and reflexivity or Arens regularity of some Banach algebras, and we investigate some applications of this new definition in the weak amenability of some Banach algebras.

1. preliminaries and Introduction

As is well-known [1], the second dual A^{**} of A endowed with the either Arens multiplications is a Banach algebra. The constructions of the two Arens multiplications in A^{**} lead us to definition of topological centers for A^{**} with respect both Arens multiplications. The topological centers of Banach algebras, module actions and applications of them were introduced and discussed in [6, 8, 13, 14, 15, 16, 17, 21, 23, 24], and they have attracted by some attentions. In this paper we will introduce a new concept as weak topological center of Banach algebras, module actions and we investigate its relationship with topological centers of Banach algebras, module actions and we will obtain some conclusions in Banach algebras with some examples in the group algebras. In final section, we give some applications of weak topological center of Banach algebras in weak amenability of Banach algebras.

Now we introduce some notations and definitions that we used throughout this paper. Let A be a Banach algebra and A^* , A^{**} , respectively, are the first and second dual of A . For $a \in A$ and $a' \in A^*$, we denote by $a'a$ and aa' respectively, the functionals on A^* defined by $\langle a'a, b \rangle = \langle a', ab \rangle = a'(ab)$ and $\langle aa', b \rangle = \langle a', ba \rangle = a'(ba)$ for all $b \in A$. The Banach algebra A is embedded in its second dual via the identification $\langle a, a' \rangle = \langle a', a \rangle$ for every $a \in A$ and $a' \in A^*$. Let X, Y, Z be normed spaces and $m : X \times Y \rightarrow Z$ be a bounded bilinear mapping. Arens in [1] offers two natural extensions m^{***} and m^{t***} of m from $X^{**} \times Y^{**}$ into Z^{**} as following

1. $m^* : Z^* \times X \rightarrow Y^*$, given by $\langle m^*(z', x), y \rangle = \langle z', m(x, y) \rangle$ where $x \in X$, $y \in Y$, $z' \in Z^*$,
2. $m^{**} : Y^{**} \times Z^* \rightarrow X^*$, given by $\langle m^{**}(y'', z'), x \rangle = \langle y'', m^*(z', x) \rangle$ where $x \in X$, $y'' \in Y^{**}$, $z' \in Z^*$,
3. $m^{***} : X^{**} \times Y^{**} \rightarrow Z^{**}$, given by $\langle m^{***}(x'', y''), z' \rangle = \langle x'', m^{**}(y'', z') \rangle$ where $x'' \in X^{**}$, $y'' \in Y^{**}$, $z' \in Z^*$.

2000 *Mathematics Subject Classification.* 46L06; 46L07; 46L10; 47L25.

Key words and phrases. Arens regularity, topological center, weak topological center, module action, n -th dual, weak amenability, derivation, group algebra.

The mapping m^{***} is the unique extension of m such that $x'' \rightarrow m^{***}(x'', y'')$ from X^{**} into Z^{**} is *weak* - weak** continuous for every $y'' \in Y^{**}$, but the mapping $y'' \rightarrow m^{***}(x'', y'')$ is not in general *weak* - weak** continuous from Y^{**} into Z^{**} unless $x'' \in X$. Hence the first topological center of m may be defined as following

$$Z_1(m) = \{x'' \in X^{**} : y'' \rightarrow m^{***}(x'', y'') \text{ is } \textit{weak*} - \textit{weak*} - \textit{continuous}\}.$$

Let now $m^t : Y \times X \rightarrow Z$ be the transpose of m defined by $m^t(y, x) = m(x, y)$ for every $x \in X$ and $y \in Y$. Then m^t is a continuous bilinear map from $Y \times X$ to Z , and so it may be extended as above to $m^{t***} : Y^{**} \times X^{**} \rightarrow Z^{**}$. The mapping $m^{t***} : X^{**} \times Y^{**} \rightarrow Z^{**}$ in general is not equal to m^{***} , see [1], if $m^{***} = m^{t***t}$, then m is called Arens regular. The mapping $y'' \rightarrow m^{t***t}(x'', y'')$ is *weak* - weak** continuous for every $y'' \in Y^{**}$, but the mapping $x'' \rightarrow m^{t***t}(x'', y'')$ from X^{**} into Z^{**} is not in general *weak* - weak** continuous for every $y'' \in Y^{**}$. So we define the second topological center of m as

$$Z_2(m) = \{y'' \in Y^{**} : x'' \rightarrow m^{t***t}(x'', y'') \text{ is } \textit{weak*} - \textit{weak*} - \textit{continuous}\}.$$

It is clear that m is Arens regular if and only if $Z_1(m) = X^{**}$ or $Z_2(m) = Y^{**}$. Arens regularity of m is equivalent to the following

$$\lim_i \lim_j \langle z', m(x_i, y_j) \rangle = \lim_j \lim_i \langle z', m(x_i, y_j) \rangle,$$

whenever both limits exist for all bounded sequences $(x_i)_i \subseteq X$, $(y_i)_i \subseteq Y$ and $z' \in Z^*$, see [6, 14, 18].

The regularity of a normed algebra A is defined to be the regularity of its algebra multiplication when considered as a bilinear mapping. Let a'' and b'' be elements of A^{**} , the second dual of A . By *Goldstine's* Theorem [4, P.424-425], there are nets $(a_\alpha)_\alpha$ and $(b_\beta)_\beta$ in A such that $a'' = \textit{weak*} - \lim_\alpha a_\alpha$ and $b'' = \textit{weak*} - \lim_\beta b_\beta$. So it is easy to see that for all $a' \in A^*$,

$$\lim_\alpha \lim_\beta \langle a', m(a_\alpha, b_\beta) \rangle = \langle a''b'', a' \rangle$$

and

$$\lim_\beta \lim_\alpha \langle a', m(a_\alpha, b_\beta) \rangle = \langle a''ob'', a' \rangle,$$

where $a''b''$ and $a''ob''$ are the first and second Arens products of A^{**} , respectively, see [6, 14, 18].

The mapping m is left strongly Arens irregular if $Z_1(m) = X$ and m is right strongly Arens irregular if $Z_2(m) = Y$.

Regarding A as a Banach A -*bimodule*, the operation $\pi : A \times A \rightarrow A$ extends to π^{***} and π^{t***t} defined on $A^{**} \times A^{**}$. These extensions are known, respectively, as the first (left) and the second (right) Arens products, and with each of them, the second dual space A^{**} becomes a Banach algebra. In this situation, we shall also simplify our notations. So the first (left) Arens product of $a'', b'' \in A^{**}$ shall be simply indicated by $a''b''$ and defined by the three steps:

$$\begin{aligned} \langle a'a, b \rangle &= \langle a', ab \rangle, \\ \langle a''a', a \rangle &= \langle a'', a'a \rangle, \\ \langle a''b'', a' \rangle &= \langle a'', b''a' \rangle. \end{aligned}$$

for every $a, b \in A$ and $a' \in A^*$. Similarly, the second (right) Arens product of $a'', b'' \in A^{**}$ shall be indicated by $a''ob''$ and defined by :

$$\begin{aligned}\langle a'oa', b \rangle &= \langle a', ba \rangle, \\ \langle a'oa'', a \rangle &= \langle a'', a'oa' \rangle, \\ \langle a''ob'', a' \rangle &= \langle b'', a'ob'' \rangle.\end{aligned}$$

for all $a, b \in A$ and $a' \in A^*$.

The regularity of a normed algebra A is defined to be the regularity of its algebra multiplication when considered as a bilinear mapping. Let a'' and b'' be elements of A^{**} , the second dual of A . By *Goldstine's* Theorem [4, P.424-425], there are nets $(a_\alpha)_\alpha$ and $(b_\beta)_\beta$ in A such that $a'' = \text{weak}^* - \lim_\alpha a_\alpha$ and $b'' = \text{weak}^* - \lim_\beta b_\beta$. So it is easy to see that for all $a' \in A^*$,

$$\lim_\alpha \lim_\beta \langle a', \pi(a_\alpha, b_\beta) \rangle = \langle a''b'', a' \rangle$$

and

$$\lim_\beta \lim_\alpha \langle a', \pi(a_\alpha, b_\beta) \rangle = \langle a''ob'', a' \rangle,$$

where $a''b''$ and $a''ob''$ are the first and second Arens products of A^{**} , respectively, see [6, 14, 18].

We find the usual first and second topological center of A^{**} , which are

$$\begin{aligned}Z_1(A^{**}) &= Z_1^\ell(A^{**}) = \{a'' \in A^{**} : b'' \rightarrow a''b'' \text{ is weak}^* - \text{weak}^* \text{ continuous}\}, \\ Z_2(A^{**}) &= Z_2^r(A^{**}) = \{a'' \in A^{**} : a'' \rightarrow a''ob'' \text{ is weak}^* - \text{weak}^* \text{ continuous}\}.\end{aligned}$$

An element e'' of A^{**} is said to be a mixed unit if e'' is a right unit for the first Arens multiplication and a left unit for the second Arens multiplication. That is, e'' is a mixed unit if and only if, for each $a'' \in A^{**}$, $a''e'' = e''oa'' = a''$. By [4, p.146], an element e'' of A^{**} is mixed unit if and only if it is a *weak*^{*} cluster point of some BAI $(e_\alpha)_{\alpha \in I}$ in A .

Let now B be a Banach A - *bimodule*, and let

$$\pi_\ell : A \times B \rightarrow B \text{ and } \pi_r : B \times A \rightarrow B.$$

be the right and left module actions of A on B . Then B^{**} is a Banach A^{**} - *bimodule* with module actions

$$\pi_\ell^{***} : A^{**} \times B^{**} \rightarrow B^{**} \text{ and } \pi_r^{***} : B^{**} \times A^{**} \rightarrow B^{**}.$$

Similarly, B^{**} is a Banach A^{**} - *bimodule* with module actions

$$\pi_\ell^{t***} : A^{**} \times B^{**} \rightarrow B^{**} \text{ and } \pi_r^{t***} : B^{**} \times A^{**} \rightarrow B^{**}.$$

Let B be a left Banach A - *module* and e be a left unit element of A . We say that e is a left unit (resp. weakly left unit) A - *module* for B , if $\pi_\ell(e, b) = b$ (resp. $\langle b', \pi_\ell(e, b) \rangle = \langle b', b \rangle$ for all $b' \in B^*$) where $b \in B$. The definition of right unit (resp. weakly right unit) A - *module* is similar.

We say that a Banach A - *bimodule* B is a unital A - *module*, if B has left and right unit A - *module* that are equal then we say that B is unital A - *module*.

Let B be a left Banach A - *module* and $(e_\alpha)_\alpha \subseteq A$ be a LAI [resp. weakly left approximate identity(=WLAI)] for A . We say that $(e_\alpha)_\alpha$ is left approximate identity

(= LAI)[resp. weakly left approximate identity (=WLAI)] for B , if for all $b \in B$, $\pi_\ell(e_\alpha, b) \rightarrow b$ (resp. $\pi_\ell(e_\alpha, b) \xrightarrow{w} b$). The definition of the right approximate identity (=RAI)[resp. weakly right approximate identity (=WRAI)] is similar.

$(e_\alpha)_\alpha \subseteq A$ is called a approximate identity (=AI)[resp. weakly approximate identity (WAI)] for B , if B has the same left and right approximate identity [resp. weakly left and right approximate identity].

Let $(e_\alpha)_\alpha \subseteq A$ be *weak** left approximate identity for A^{**} . We say that $(e_\alpha)_\alpha$ is *weak** left approximate identity as A^{**} -module (= W^*LAI as A^{**} -module) for B^{**} , if for all $b'' \in B^{**}$, we have $\pi_\ell^{***}(e_\alpha, b'') \xrightarrow{w^*} b''$. The definition of the *weak** right approximate identity (= W^*RAI) is similar.

$(e_\alpha)_\alpha \subseteq A$ is called a *weak** approximate identity (= W^*AI) for B^{**} , if B^{**} has the same *weak** left and right approximate identity.

In many parts of this paper, for left or right Banach A -module, we take $\pi_\ell(a, b) = ab$ and $\pi_r(b, a) = ba$, for every $a \in A$ and $b \in B$.

This paper is organized as follows:

- a) Suppose that A is a Banach algebra and $A^{***}A^{**} \subseteq A^*$. Then the topological centers of A^{**} ($=Z_1(A^{**})$) and weak topological centers of A^{**} ($=\tilde{Z}_1^\ell(A^{**})$) are A^{**} .
- b) Let A be a Banach algebra and let the left weak topological center of A^{**} be A . Then A is a right ideal in A^{**} .
- c) For a Banach algebra A , we have the following assertions: i) If $A^{***}A^{**} \subseteq \widehat{wap}(A)$, then $A^{**}A^{**} \subseteq Z_1(A^{**})$ where

$$\widehat{wap}(A) = \{a''' \in A^{***} : a'''a'' : A^{**} \rightarrow \mathbb{C} \text{ is weak* continuous for all } a'' \in A^{**}\}.$$

- ii) If $A^{**} = A^{**}\tilde{Z}_1^\ell(A^{**})$, then $A^{***}A^{**} \subseteq \widehat{wap}(A)$.

- d) For a compact group G , we have .

$$L^1(G) = \tilde{Z}_{M(G)^{**}}^\ell(L^1(G)^{**}), \text{ and } L^1(G) = \tilde{Z}_{M(G)^{**}}^r(L^1(G)^{**}).$$

- e) Let B a right Banach A -module and $e^{(n)} \in A^{(n)}$ be a right unit for $B^{(n)}$ where $n \geq 2$. Then we have the following assertions.

- i) If $Z_{e^{(n)}}^\ell(B^{(n)}) = B^{(n)}$, then $A^{(n-2)}B^{(n-1)} = B^{(n-1)}$ where

$$\tilde{Z}_{e^{(n)}}^\ell(B^{(n)}) = \{b^{(n)} \in B^{(n)} : e'' \rightarrow b^{(n)}e^{(n)} \text{ is weak* - to - weak continuous}\},$$

- ii) If $\tilde{Z}_{e^{(n)}}^\ell(B^{(n)}) = B^{(n)}$, then $A^{(n-2)}B^{(n-1)} = B^{(n-1)}$ and *weak** closure of $A^{(n-2)}B^{(n+1)}$ is $B^{(n+1)}$.

- f) Assume that A is a Banach algebra and $\tilde{Z}_1^\ell(A^{(n)}) = A^{(n)}$ where $n > 1$. If $A^{(n)}$ is weakly amenable, then $A^{(n-2)}$ is weakly amenable.

- g) Let B a Banach A -bimodule and $D : A^{(n)} \rightarrow B^{(n+1)}$ be a derivation where $n \geq 0$. If $\tilde{Z}_{A^{(n)}}^\ell(B^{(n)}) = B^{(n)}$, then $D'' : A^{(n+2)} \rightarrow A^{(n+3)}$ is a derivation.

- j) Let B be a Banach A -bimodule and $\tilde{Z}_{A^{**}}^\ell(B^{**}) = B^{**}$. Assume that $D : A \rightarrow B^*$ is a derivation. Then $D^{**}(A^{**})B^{**} \subseteq A^*$.

2. Weak topological center of Banach algebras

In this section, we define a new concept as weak topological center of the second dual of a Banach algebra A , A^{**} , with respect to first and second Arens product and

we find its relationships with the topological centers of A^{**} with respect to first and second Arens products. We establish some conditions that when the weak topological center and topological center of A^{**} coincide.

Definition 2-1. Let A be a Banach algebra. Then we define weak topological centers of A^{**} with respect to the first and second Arens product, respectively, in the following

$$\begin{aligned}\tilde{Z}_1^\ell(A^{**}) &= \{a'' \in A^{**} : b'' \rightarrow a''b'' \text{ is weak}^* - \text{to} - \text{weak continuous in } A^{**}\}, \\ \tilde{Z}_1^r(A^{**}) &= \{a'' \in A^{**} : b'' \rightarrow b''a'' \text{ is weak}^* - \text{to} - \text{weak continuous in } A^{**}\}, \\ \tilde{Z}_2^\ell(A^{**}) &= \{a'' \in A^{**} : b'' \rightarrow b''oa'' \text{ is weak}^* - \text{to} - \text{weak continuous in } A^{**}\}, \\ \tilde{Z}_2^r(A^{**}) &= \{a'' \in A^{**} : b'' \rightarrow a''ob'' \text{ is weak}^* - \text{to} - \text{weak continuous in } A^{**}\}.\end{aligned}$$

It is clear that $\tilde{Z}_i^\ell(A^{**})$ and $\tilde{Z}_i^r(A^{**})$ for each $i \in \{1, 2\}$, are subspace of A^{**} with respect to the first and second Arens products. In general, we have the following results:

- (1) $\tilde{Z}_1^\ell(A^{**}) \subseteq Z_1(A^{**})$, $\tilde{Z}_2^r(A^{**}) \subseteq Z_2(A^{**})$.
- (2) If $Z_1(A^{**}) = Z_2(A^{**})$, it follow that $\tilde{Z}_1^\ell(A^{**}) = \tilde{Z}_2^\ell(A^{**})$ and $\tilde{Z}_1^r(A^{**}) = \tilde{Z}_2^r(A^{**})$.
- (3) If $Z_1(A^{**}) = \tilde{Z}_1^\ell(A^{**}) = \tilde{Z}_2^r(A^{**})$ or $\tilde{Z}_1^\ell(A^{**}) = \tilde{Z}_2^r(A^{**}) = Z_2(A^{**})$, then we conclude that $\tilde{Z}_1(A^{**}) = Z_2(A^{**})$.
- (4) If $\tilde{Z}_1^\ell(A^{**}) = A^{**}$ or $\tilde{Z}_1^r(A^{**}) = A^{**}$, then A is Arens regular.
- (5) Let $\tilde{Z}_1^\ell(A^{**}) = Z_1(A^{**})$. Then for every subspace B of A^{**} , we have $B\tilde{Z}_1^\ell(A^{**}) = BZ_1(A^{**})$.

If a Banach algebra A is Arens regular or strongly Arens irregular on the left, in general, we can not conclude that $\tilde{Z}_1^\ell(A^{**}) = A^{**}$ or $\tilde{Z}_1^\ell(A^{**}) = A$, respectively. In the following, we give examples of some Arens regular or strongly Arens irregular Banach algebras such as A that $\tilde{Z}_1^\ell(A^{**})$ is equal to A^{**} or no.

- i) Let A be nonreflexive Arens regular Banach algebra and $e'' \in A^{**}$ be a left unite element of A^{**} . Then $\tilde{Z}_1^\ell(A^{**}) \neq A^{**}$.
- ii) Suppose that G is a locally compact group. Then by notice to [17], we know that in spacial case, $M(G)$ is left strong Arens irregular, but $\tilde{Z}_1^\ell(M(G)^{**}) \neq M(G)$. Of course, it is clear that $\tilde{Z}_1^\ell(M(G)^{**}) \subsetneq M(G)$.

Now in the following we give an example nonreflexive Arnse regular Banach algebra which $\tilde{Z}_1^\ell(A^{**}) = A^{**}$.

Example 2-2. Consider the algebra $c_0 = (c_0, \cdot)$ is the collection of all sequences of scalars that convergence to 0, with the some vector space operations and norm as ℓ_∞ . We show that $\tilde{Z}_1^\ell(c_0^{**}) = c_0^{**}$.

Proof. We know that $c_0^* = \ell^1$, $c_0^{**} = \ell^\infty$ and $c_0^{***} = (\ell^\infty)^* = \ell^{1**} = \ell^1 \oplus c_0^\perp$ as a Banach c_0 - bimodule. Take $a''' \in c_0^{***}$. Then $a''' = (a', t)$ where $a' \in \ell^1$ and $t \in c_0^\perp$. By [6, 2.6.22], $a''a' \in c_0^\perp$ for all $a'' \in c_0^{**} = \ell^\infty$. Then we have

$$\langle ta'', a' \rangle = \langle t, a''a' \rangle = 0.$$

It follows that $ta'' = 0$. So for all $b'' \in c_0^{**} = \ell^\infty$, we have the following equalities

$$\begin{aligned}\langle a''', a''b'' \rangle &= \langle a', a''b'' \rangle \oplus \langle t, a''b'' \rangle = \langle a''b'', a' \rangle \oplus \langle ta'', b'' \rangle \\ &= \langle a''b'', a' \rangle.\end{aligned}$$

Consequently $\tilde{Z}_1^\ell(c_0^{**}) = Z_1(c_0^{**})$. Since c_0^{**} is Arens regular, $\tilde{Z}_1^\ell(c_0^{**}) = c_0^{**}$. $\square$

Theorem 2-3. Suppose that A is a Banach algebra and $B \subseteq A^{**}$. Then we have the following statements.

- (1) If $A^{***}B \subseteq A^*$, then $B \subseteq \tilde{Z}_1^\ell(A^{**})$.
- (2) If $BA^{***} \subseteq A^*$, then $B \subseteq \tilde{Z}_1^r(A^{**})$.

Proof. 1) Let $b \in B$ and $(a''_\alpha)_\alpha \subseteq A^{**}$ such that $a''_\alpha \xrightarrow{w^*} a''$. We shows that $ba''_\alpha \xrightarrow{w} ba''$. Let $a''' \in A^{***}$. Since $A^{***}B \subseteq A^*$, $a'''b \in A^*$. Then we have

$$\langle a''', ba''_\alpha \rangle = \langle a'''b, a''_\alpha \rangle = \langle a''_\alpha, a'''b \rangle \rightarrow \langle a''_\alpha, a'''b \rangle = \langle a''_\alpha, a'''b \rangle = \langle a''', ba'' \rangle.$$

2) Proof is similar to (1). $\square$

Corollary 2-4. Suppose that A is a Banach algebra. Then we have the following statements.

- (1) If $A^{***}A^{**} \subseteq A^*$, then $\tilde{Z}_1^\ell(A^{**}) = Z_1(A^{**}) = A^{**}$.
- (2) If $A^{**}A^{***} \subseteq A^*$, then $\tilde{Z}_1^r(A^{**}) = A^{**}$.

Corollary 2-5. Suppose that A is a Banach algebra and $A^{***}A \subseteq A^*$. If A is left strong Arens irregular, then $\tilde{Z}_1^\ell(A^{**}) = A$.

Lemma 2-6. Suppose that A is a Banach algebra. Let $(x''_\alpha)_\alpha \subseteq A^{**}$ and $(y'_\beta)_\beta \subseteq A^*$ have taken *weak** limit in A^{**} and A^{***} , respectively. Let

$$\lim_{\beta} \lim_{\alpha} \langle x''_\alpha, y'_\beta \rangle = \lim_{\alpha} \lim_{\beta} \langle x''_\alpha, y'_\beta \rangle,$$

then we have the following assertions.

- (1) $\tilde{Z}_1^r(A^{**}) = A^{**}$.
- (2) $\tilde{Z}_1^\ell(A^{**}) = Z_1(A^{**})$.

Proof. 1) Let $b'' \in A^{**}$. For all $a''' \in A^{***}$ there is $(a'_\beta)_\beta \subseteq A^*$ such that $a'_\beta \xrightarrow{w^*} a'''$.

Assume that $(a''_\alpha)_\alpha \subseteq A^{**}$ such that $a''_\alpha \xrightarrow{w^*} a''$. Then we have

$$\begin{aligned}\langle a''', a''b'' \rangle &= \lim_{\beta} \langle a'_\beta, a''b'' \rangle = \lim_{\beta} \lim_{\alpha} \langle a'_\beta, a''_\alpha b'' \rangle = \lim_{\alpha} \lim_{\beta} \langle a'_\beta, a''_\alpha b'' \rangle \\ &= \lim_{\alpha} \langle a''', a''_\alpha b'' \rangle.\end{aligned}$$

It follows that the mapping $a'' \rightarrow a''b''$ is *weak* - to - weak continuous* for all $b'' \in A^{**}$, and so $\tilde{Z}_1^r(A^{**}) = A^{**}$.

2) Proof is similar to (1). $\square$

By conditions Lemma 2-6, it is clear that if A is Arens regular, then we have $\tilde{Z}_1^\ell(A^{**}) = \tilde{Z}_1^r(A^{**}) = A^{**}$.

Theorem 2-7. Let A be a Banach algebra and let $B \subseteq A^{**}$. Then we have $B\tilde{Z}_1^\ell(A^{**}) \subseteq \tilde{Z}_1^\ell(A^{**})$ and $\tilde{Z}_1^r(A^{**})B \subseteq \tilde{Z}_1^r(A^{**})$.

Proof. We prove that $B\tilde{Z}_1^\ell(A^{**}) \subseteq \tilde{Z}_1^\ell(A^{**})$ and proof of the other part is the same. Suppose that $a'' \in \tilde{Z}_1^\ell(A^{**})$ and $b'' \in B$. Take $a''' \in A^{***}$ and $(c''_\alpha)_\alpha \subseteq A^{**}$ such that $c''_\alpha \xrightarrow{w^*} c''$. Then

$$\langle a''', b''a''c''_\alpha \rangle = \langle a'''b'', a''c''_\alpha \rangle \rightarrow \langle a'''b'', a''c'' \rangle = \langle a''', b''a''c'' \rangle.$$

It follows that the mapping $c'' \rightarrow (b''a'')c''$ is *weak* - to - weak* continuous, and so $b''a'' \in \tilde{Z}_1^\ell(A^{**})$. $\square$

Corollary 2-8. Let A be a Banach algebra. Then we have the following assertions.

- (1) If $\tilde{Z}_1^\ell(A^{**}) = A$, then A is a right ideal in A^{**} .
- (2) If $\tilde{Z}_2^\ell(A^{**}) = A$, then A is a left ideal in A^{**} .
- (3) If $\tilde{Z}_1^\ell(A^{**}) = \tilde{Z}_2^\ell(A^{**}) = A$, then A is an ideal in A^{**} .

Let A be a Banach algebra. The subspace of A^{***} annihilating A will be denoted by $A^\perp = \{a''' \in A^{***} \mid a'''|_A = 0\}$.

Theorem 2-9. Let A be a Banach algebra. Then we have the following assertions.

- (1) If A is a left ideal in A^{**} , then $A \subseteq \tilde{Z}_1^\ell(A^{**})$.
- (2) If A is a right ideal in A^{**} , then $A \subseteq \tilde{Z}_2^\ell(A^{**})$.
- (3) If A is an ideal in A^{**} , then $A \subseteq \tilde{Z}_1^\ell(A^{**}) \cap \tilde{Z}_2^\ell(A^{**})$.
- (4) If A is a left (resp. right) ideal in A^{**} and A^{**} has a left (resp. right) unit element that is belong to $\tilde{Z}_1^\ell(A^{**})$ (resp. $\tilde{Z}_2^\ell(A^{**})$). Then A is reflexive.

Proof. (1) Assume that $(a''_\alpha)_\alpha \subseteq A^{**}$ such that $a''_\alpha \xrightarrow{w^*} a''$. Let $a''' \in A^{***}$. Since $A^{***} = A^* \oplus A^\perp$, there are $a' \in A^*$ and $t \in A^\perp$ such that $a''' = (a', t)$. Then for every $a \in A$, we have

$$\begin{aligned} \langle a''', aa''_\alpha \rangle &= \langle (a', t), aa''_\alpha \rangle = \langle aa''_\alpha, a' \rangle \\ &\rightarrow \langle aa'', a' \rangle = \langle a''', aa'' \rangle. \end{aligned}$$

It follows that $A \subseteq \tilde{Z}_1^\ell(A^{**})$.

- (2) Proof is similar to (1).
- (3) Proof is clear.
- (4) Let $e \in \tilde{Z}_1^\ell(A^{**})$ be a unit element for A^{**} . Take $(a''_\alpha)_\alpha \subseteq A^{**}$ such that $a''_\alpha \xrightarrow{w^*} a''$ and $a''' \in A^{***}$. Since $A^{***} = A^* \oplus A^\perp$, there are $a' \in A^*$ and $t \in A^\perp$ such that $a''' = (a', t)$. Thus

$$\begin{aligned} \langle a''', a''_\alpha \rangle &= \langle a''', ea''_\alpha \rangle = \langle (a', t), ea''_\alpha \rangle = \langle ea''_\alpha, a' \rangle \\ &\rightarrow \langle ea'', a' \rangle = \langle a''', a'' \rangle. \end{aligned}$$

It follows that $a''_\alpha \xrightarrow{w} a''$. Hence A is reflexive. $\square$

Corollary 2-10. Let A be a Banach algebra. Then we have the following assertions.

- (1) If A is a left ideal in A^{**} and A left strongly Arens irregular, then $A = \tilde{Z}_1^\ell(A^{**})$, and so A is an ideal in A^{**} .
- (2) If A is a right ideal in A^{**} and A right strongly Arens irregular, then $A = \tilde{Z}_2^\ell(A^{**})$, and so A is an ideal in A^{**} .
- (3) If A is an ideal in A^{**} and A strongly Arens irregular, then $A = \tilde{Z}_1^\ell(A^{**}) = \tilde{Z}_2^\ell(A^{**})$.
- (4) If A is a left (resp. right) ideal in A^{**} and A^{**} has a left (resp. right) unit element belong to $\tilde{Z}_1^\ell(A^{**})$ (resp. $\tilde{Z}_2^\ell(A^{**})$). Then A is reflexive.

Example 2-11. i) Let G be a compact group. By using [13], we know that $L^1(G)$ is a strongly Arens irregular and by [19], $L^1(G)$ is an ideal in its second dual. Then $L^1(G) = \tilde{Z}_1^\ell(L^1(G)^{**}) = \tilde{Z}_2^\ell(L^1(G)^{**})$.

ii) Let G be a locally compact group. Then, in general, $M(G)$ is not a left or right ideal in its second dual.

iii) Let G be a locally compact group. If $L^1(G)$ is a left or right ideal in its second dual, it is an ideal in its second dual.

A functional a' in A^* is said to be *wap* (weakly almost periodic) on A if the mapping $a \rightarrow a'a$ from A into A^* is weakly compact. Pym in [18] showed that this definition to the equivalent following condition

For any two net $(a_\alpha)_\alpha$ and $(b_\beta)_\beta$ in $\{a \in A : \|a\| \leq 1\}$, we have

$$\lim_\alpha \lim_\beta \langle a', a_\alpha b_\beta \rangle = \lim_\beta \lim_\alpha \langle a', a_\alpha b_\beta \rangle,$$

whenever both iterated limits exist. The collection of all *wap* functionals on A is denoted by $wap(A)$. Also we have $a' \in wap(A)$ if and only if $\langle a''b'', a' \rangle = \langle a''ob'', a' \rangle$ for every $a'', b'' \in A^{**}$.

Definition 2-12. Let A be a Banach algebra. We define $\widetilde{wap}(A)$ as a subset of A^{***} as follows

$$\widetilde{wap}_\ell(A) = \{a''' \in A^{***} : a'''a'' : A^{**} \rightarrow \mathbb{C} \text{ is weak}^* \text{ continuous for all } a'' \in A^{**}\}.$$

It is clear that $\widetilde{wap}_\ell(A) = A^{***}$ if and only if $\tilde{Z}_1^\ell(A^{**}) = A^{**}$. Thus, if $\widetilde{wap}_\ell(A) = A^{***}$, then $wap(A) = A^*$, and so A is Arens regular. It is easy to show that $wap(A) = \widetilde{wap}_\ell(A)$ if and only if $\tilde{Z}_1^\ell(A^{**}) = Z_1(A^{**})$.

By notice to Examples 2-2, it is clear that $\widetilde{wap}_\ell(c_0) = c_0^{***}$, but for locally compact group G , we have $\widetilde{wap}_\ell(L^1(G)) \neq L^1(G)^{***}$.

The definition of $\widetilde{wap}_r(A)$ is similar.

Theorem 2-13. Let A be a Banach algebra. Then we have the following assertions.

- (1) If $A^{***}A^{**} \subseteq \widehat{wap}_\ell(A)$, then $\tilde{Z}_1^\ell(A^{**}) = Z_1(A^{**}) = A^{**}$.
- (2) If $A^{**} = A^{**}\tilde{Z}_1^\ell(A^{**})$, then $A^{***}A^{**} \subseteq \widehat{wap}_\ell(A)$.

Proof. 1) Suppose that $a'' \in A^{**}$ and $(b''_\alpha)_\alpha \subseteq A^{**}$ such that $b''_\alpha \xrightarrow{w^*} b''$. Take $a''' \in A^{***}$. Then, since $A^{***}A^{**} \subseteq \widehat{wap}_\ell(A)$, we have

$$\langle a''', a''b''_\alpha \rangle = \langle a'''a'', b''_\alpha \rangle \rightarrow \langle a'''a'', b'' \rangle = \langle a''', a''b'' \rangle.$$

It follows that $a'' \in \tilde{Z}_1^\ell(A^{**})$, and so $A^{**} = \tilde{Z}_1^\ell(A^{**})$. Since $\tilde{Z}_1^\ell(A^{**}) \subseteq Z_1(A^{**})$, the result is hold.

2) Let $a'' \in A^{**}$ and $a''' \in A^{***}$. Since $A^{**} = A^{**}\tilde{Z}_1^\ell(A^{**})$, there are $b'' \in A^{**}$ and $c'' \in \tilde{Z}_1^\ell(A^{**})$ such that $a'' = b''c''$. Consequently, we have

$$\langle a'''a'', d'' \rangle = \langle a'''b'', c''d'' \rangle,$$

where $d'' \in A^{**}$. Since $c'' \in \tilde{Z}_1^\ell(A^{**})$, $a''' \in \widehat{wap}_\ell(A)$. □

Corollary 2-14. Let A be a Banach algebra. If $A^{**} = A^{**}\tilde{Z}_1^\ell(A^{**})$, then $\tilde{Z}_1^\ell(A^{**}) = Z_1(A^{**}) = A^{**}$.

Proof. Since $wap(A) \subseteq \widehat{wap}_\ell(A)$, by using preceding theorem, proof is hold. □

Corollary 2-15. Let A be a Banach algebra. Then, if $A^{***}A^{**} \subseteq wap(A)$, then $\tilde{Z}_1^\ell(A^{**}) = Z_1(A^{**}) = A^{**}$.

Corollary 2-16. Let A be a Banach algebra and B be a subspace of A^{**} . If $B = B\tilde{Z}_1^\ell(B)$, then $\tilde{Z}_1^\ell(B) = Z_1(B) = B$.

3. Weak topological center of module actions

In this section, we study the definition of the weak topological centers on module actions and establish some relations between it and reflexivity (spacial Arens regularity) of Banach algebras. For a Banach A – bimodule B we will study this definition on n – th dual of A and B . In some parts of this section, we have some conclusions in the group algebras, that is, for a locally compact group G and a mixed unit e'' of $L^1(G)^{**}$, we have $Z_{e''}^\ell(L^1(G)^{**}) \neq L^1(G)^{**}$ and $Z_{e''}^r(L^1(G)^{**}) \neq L^1(G)^{**}$ where $Z_{e''}^\ell(L^1(G)^{**})$ is a locally topological center of $L^1(G)^{**}$.

We have also conclusions as $\tilde{Z}_{L^1(G)^{**}}^\ell(L^1(G)^{***}) \subseteq L^1(G)^*$ and $\tilde{Z}_{L^1(G)^{***}}^r(L^1(G)^{**}) \subseteq L^1(G)$. On the other hand for Banach algebras c_0 and its second dual ℓ^∞ , we conclude that $\tilde{Z}_{\ell^\infty}^\ell((\ell^\infty)^*) \subseteq \ell^\infty$ and $\tilde{Z}_{(\ell^\infty)^*}^r(\ell^\infty) \subseteq c_0$ where multiplications in c_0^{**} and ℓ^∞

coincide.

We also extend some results from [8] into general situations and for a locally compact group G and $n \geq 3$, we have the following equalities

- (1) $Z_{L^1(G)^{(n)}}^\ell(L^1(G)^{**}) = L^1(G)$, see [8].
- (2) $Z_{M(G)^{(n)}}^\ell(L^1(G)^{**}) = L^1(G)$.
- (3) $Z_{M(G)^{(n)}}^\ell(M(G)^{**}) = M(G)$.
- (4) $Z_{L^1(G)^{**}}^\ell(L^1(G)^{***}) = L^1(G)^*$ (resp. $Z_{L^1(G)^{**}}^r(L^1(G)^{***}) = L^1(G)^*$) where multiplication is to the first (resp. second) Arens product.
- (5) Let $n > 0$ be even number. Then $Z_{M(G)^{(n)}}^\ell(M(G)^{(n+1)}) = M(G)^{(n-1)}$ (resp. $Z_{M(G)^{(n)}}^r(M(G)^{(n+1)}) = M(G)^{(n-1)}$) where multiplication is to the first (resp. second) Arens product.
- (6) For a compact group G , we have

$$L^1(G) = \tilde{Z}_{M(G)^{**}}^\ell(L^1(G)^{**}) = \tilde{Z}_{M(G)^{**}}^r(L^1(G)^{**}).$$

Suppose that A is a Banach algebra and B is a Banach A – *bimodule*. According to [5, pp.27 and 28], B^{**} is a Banach A^{**} – *bimodule*, where A^{**} is equipped with the first Arens product. So we define the topological centers and weak topological centers of module actions. First for a Banach A – *bimodule* B , we define the topological centers of the left and right module actions of A on B as follows:

$$\begin{aligned} Z_{A^{**}}^\ell(B^{**}) &= Z(\pi_r) = \{b'' \in B^{**} : \text{the map } a'' \rightarrow \pi_r^{***}(b'', a'') : A^{**} \rightarrow B^{**} \\ &\quad \text{is weak}^* - \text{weak}^* \text{ continuous}\} \\ Z_{B^{**}}^\ell(A^{**}) &= Z(\pi_\ell) = \{a'' \in A^{**} : \text{the map } b'' \rightarrow \pi_\ell^{***}(a'', b'') : B^{**} \rightarrow B^{**} \\ &\quad \text{is weak}^* - \text{weak}^* \text{ continuous}\} \\ Z_{A^{**}}^r(B^{**}) &= Z(\pi_\ell^t) = \{b'' \in B^{**} : \text{the map } a'' \rightarrow \pi_\ell^{t***}(b'', a'') : A^{**} \rightarrow B^{**} \\ &\quad \text{is weak}^* - \text{weak}^* \text{ continuous}\} \\ Z_{B^{**}}^r(A^{**}) &= Z(\pi_r^t) = \{a'' \in A^{**} : \text{the map } b'' \rightarrow \pi_r^{t***}(a'', b'') : B^{**} \rightarrow B^{**} \\ &\quad \text{is weak}^* - \text{weak}^* \text{ continuous}\}. \end{aligned}$$

Now we define the weak topological centers of the right and left module actions of A on B as follows:

Definition 3-1. Let B be a Banach A – *bimodule*. Then we define the weak topological centers of left and right module actions as follows

$$\begin{aligned} \tilde{Z}_{A^{**}}^\ell(B^{**}) &= \{b'' \in B^{**} : \text{the mapping } a'' \rightarrow b''a'' \text{ is weak}^* - \text{weak continuous}\}, \\ \tilde{Z}_{A^{**}}^r(B^{**}) &= \{b'' \in B^{**} : \text{the mapping } a'' \rightarrow a''b'' \text{ is weak}^* - \text{weak continuous}\}, \\ \tilde{Z}_{B^{**}}^\ell(A^{**}) &= \{a'' \in A^{**} : \text{the mapping } a'' \rightarrow b''a'' \text{ is weak}^* - \text{weak continuous}\}, \\ \tilde{Z}_{B^{**}}^r(A^{**}) &= \{a'' \in A^{**} : \text{the mapping } a'' \rightarrow a''b'' \text{ is weak}^* - \text{weak continuous}\}. \end{aligned}$$

Let $A^{(n)}$ and $B^{(n)}$ be n – *th dual* of B and A , respectively where $n \geq 0$ is even number. By [25], $B^{(n)}$ is a Banach $A^{(n)}$ – *bimodule* and we may therefore define the

topological centers and weak topological centers of the right and left module action of $A^{(n)}$ on $B^{(n)}$ to similar above. Then we define $B^{(n)}B^{(n-1)}$ as a subspace of $A^{(n-1)}$, that is, for every $b^{(n)} \in B^{(n)}$, $b^{(n-1)} \in B^{(n-1)}$ and $a^{(n-2)} \in A^{(n-2)}$, we have

$$\langle b^{(n)}b^{(n-1)}, a^{(n-2)} \rangle = \langle b^{(n)}, b^{(n-1)}a^{(n-2)} \rangle.$$

If n is odd number, we define $B^{(n)}B^{(n-1)}$ as a subspace of $A^{(n)}$, that is, for all $b^{(n)} \in B^{(n)}$, $b^{(n-1)} \in B^{(n-1)}$ and $a^{(n-1)} \in A^{(n-1)}$, we define

$$\langle b^{(n)}b^{(n-1)}, a^{(n-1)} \rangle = \langle b^{(n)}, b^{(n-1)}a^{(n-1)} \rangle;$$

and if $n = 0$, we take $A^{(0)} = A$ and $B^{(0)} = B$.

Theorem 3-2. Let B be a Banach A -bimodule. Then for every even number $n \geq 2$, we have the following assertions.

- (1) $Z_{A^{(n)}}^\ell(B^{(n+1)}) = B^{(n+1)}$ if and only if $\tilde{Z}_{A^{(n)}}^r(B^{(n)}) = B^{(n)}$.
- (2) $Z_{A^{(n)}}^\ell(A^{(n+1)}) = A^{(n+1)}$ if and only if $\tilde{Z}_1^r(A^{(n)}) = A^{(n)}$.
- (3) $Z_{A^{(n)}}^r(A^{(n+1)}) = A^{(n+1)}$ if and only if $\tilde{Z}_1^\ell(A^{(n)}) = A^{(n)}$.
- (4) $Z_{B^{(n)}}^r(A^{(n+1)}) = A^{(n+1)}$ if and only if $\tilde{Z}_{A^{(n)}}^\ell(B^{(n)}) = B^{(n)}$.

Proof. 1) Suppose that $Z_{A^{(n)}}^\ell(B^{(n+1)}) = B^{(n+1)}$ and $b^{(n)} \in B^{(n)}$. We show that the mapping $a^{(n)} \rightarrow a^{(n)}b^{(n)}$ is *weak*-to-weak* continuous. Assume that $(a_\alpha^{(n)})_\alpha \subseteq A^{(n)}$ such that $a_\alpha^{(n)} \xrightarrow{w^*} a^{(n)}$. Then for all $b^{(n+1)} \in B^{(n+1)}$, we have $b^{(n+1)}a_\alpha^{(n)} \xrightarrow{w^*} b^{(n+1)}a^{(n)}$. It follows that

$$\begin{aligned} \langle b^{(n+1)}, a_\alpha^{(n)}b^{(n)} \rangle &= \langle b^{(n+1)}a_\alpha^{(n)}, b^{(n)} \rangle \rightarrow \langle b^{(n+1)}a^{(n)}, b^{(n)} \rangle \\ &= \langle b^{(n+1)}, a^{(n)}b^{(n)} \rangle. \end{aligned}$$

Thus we conclude that $b^{(n)} \in \tilde{Z}_{A^{(n)}}^r(B^{(n)})$.

The converse is similarly.

Proof of (2), (3), (4) is similar to (1). □

In the following example, for some non-reflexive Banach algebras such as A , we show that it is maybe that $\tilde{Z}_{A^{**}}^r(A^{**})$ is equal to A^{**} or no.

Example 3-3. i) Let A be non-reflexive Banach space and let $\langle f, x \rangle = 1$ for some $f \in A^*$ and $x \in A$. We define the product on A by $ab = \langle f, b \rangle a$. It is clear that A is a Banach algebra with this product and it has right identity x , see [5]. By easy calculation, for all $a' \in A^*$, $a'' \in A^{**}$ and $a''' \in A^{***}$, we have

$$\begin{aligned} a'a &= \langle a', a \rangle f, \\ a''a' &= \langle a'', f \rangle a', \\ a'''a'' &= \langle a'', a'' \rangle \langle \cdot, f \rangle. \end{aligned}$$

Therefore we have $Z^\ell_{A^{**}}(A^{***}) \neq A^{***}$. So by Theorem 3-2, we have $\tilde{Z}^r_{A^{**}}(A^{**}) \neq A^{**}$.

Similarly, if we define the product on A as $ab = \langle f, a \rangle b$ for all $a, b \in A$, then we have $Z^\ell_{A^{**}}(A^{***}) = A^{***}$. By using Theorem 3-2, it follows that $\tilde{Z}^r_{A^{**}}(A^{**}) = A^{**}$.

ii) By using Example 2-2, we know that $\tilde{Z}^\ell_1(c_0^{**}) = c_0^{**}$, and so by Corollary 3-2, we have $Z^\ell_{c_0^{**}}(c_0^{***}) = c_0^{***}$.

Theorem 3-4. Let B be a Banach A -bimodule. Then for every odd number $n \geq 2$, we have the following assertions.

- (1) If $B^{(n)}B^{(n-1)} \subseteq A^{(n-2)}$, then $Z^\ell_{A^{(n-1)}}(B^{(n-1)}) = \tilde{Z}^\ell_{A^{(n-1)}}(B^{(n-1)})$.
- (2) If $B^{(n)}A^{(n-1)} \subseteq B^{(n-2)}$ and $B^{(n-1)}$ has a left unite $A^{(n-1)}$ -module, then B is reflexive.

Proof. 1) It is clear that $\tilde{Z}^\ell_{A^{(n-1)}}(B^{(n-1)}) \subseteq Z^\ell_{A^{(n-1)}}(B^{(n-1)})$. We show that for all $b^{(n-1)} \in Z^\ell_{A^{(n-1)}}(B^{(n-1)})$, the mapping $a^{(n-1)} \rightarrow b^{(n-1)}a^{(n-1)}$ is *weak**-to-*weak* continuous. Suppose that $(a_\alpha^{(n-1)})_\alpha \subseteq A^{(n-1)}$ and $a_\alpha^{(n-1)} \xrightarrow{w^*} a^{(n-1)}$. Since $b^{(n-1)} \in Z^\ell_{A^{(n-1)}}(B^{(n-1)})$, $b^{(n-1)}a_\alpha^{(n-1)} \xrightarrow{w^*} b^{(n-1)}a^{(n-1)}$. Take $b^{(n)} \in B^{(n)}$. Then, since $B^{(n)}B^{(n-1)} \subseteq A^{(n-2)}$, we have

$$\begin{aligned} \langle b^{(n)}, b^{(n-1)}a_\alpha^{(n-1)} \rangle &= \langle b^{(n)}b^{(n-1)}, a_\alpha^{(n-1)} \rangle = \langle a_\alpha^{(n-1)}, b^{(n)}b^{(n-1)} \rangle \\ &\rightarrow \langle a^{(n-1)}, b^{(n)}b^{(n-1)} \rangle = \langle b^{(n)}b^{(n-1)}, a^{(n-1)} \rangle = \langle b^{(n)}, b^{(n-1)}a^{(n-1)} \rangle. \end{aligned}$$

2) Let $e^{(n-1)}$ be a left unit $A^{(n-1)}$ -module for $B^{(n-1)}$ and let $(b_\alpha^{(n-1)})_\alpha \subseteq B^{(n-1)}$ such that $b_\alpha^{(n-1)} \xrightarrow{w^*} b^{(n-1)}$. Suppose that $b^{(n)} \in B^{(n)}$. Since $b^{(n)}e^{(n-1)} \in B^{(n-2)}$, we have

$$\begin{aligned} \langle b^{(n)}, b_\alpha^{(n-1)} \rangle &= \langle b^{(n)}, e^{(n-1)}b_\alpha^{(n-1)} \rangle = \langle b^{(n)}e^{(n-1)}, b_\alpha^{(n-1)} \rangle \\ &= \langle b_\alpha^{(n-1)}, b^{(n)}e^{(n-1)} \rangle \rightarrow \langle b^{(n-1)}, b^{(n)}e^{(n-1)} \rangle = \langle b^{(n-1)}, b^{(n)} \rangle. \end{aligned}$$

Consequently, the weak topology and *weak** topology on $B^{(n-1)}$ is coincide, and so $B^{(n-1)}$ is reflexive which implies that B is reflexive. $\square$

Corollary 3-5. Suppose that A is a Banach algebra. Then, for every odd number $n \geq 3$, we have the following assertions.

- (1) If $A^{(n)}A^{(n-1)} \subseteq A^{(n-2)}$, then $Z^\ell_1(A^{(n-1)}) = \tilde{Z}^\ell_1(A^{(n-1)})$.
- (2) If $A^{(n)}A^{(n-1)} \subseteq A^{(n-2)}$ and $A^{(n-3)}$ has a bounded left approximate identity ($= BLAI$), then A is reflexive.

Theorem 3-6. Let $n > 0$ be an even number and let B be a left (resp. right) Banach A -module such that $A^{(n-2)}B^{(n)} \subseteq B^{(n-2)}$ (resp. $B^{(n)}A^{(n-2)} \subseteq B^{(n-2)}$).

- (1) Then $A^{(n-2)} \subseteq \tilde{Z}^\ell_{B^{(n)}}(A^{(n)})$ (resp. $A^{(n-2)} \subseteq \tilde{Z}^r_{B^{(n)}}(A^{(n)})$).
- (2) If $B^{(n)}$ has a left (resp. right) unit element that it is belong to $\tilde{Z}^\ell_{B^{(n)}}(A^{(n)})$ (resp. $\tilde{Z}^r_{B^{(n)}}(A^{(n)})$). Then A is reflexive.

(3) If $A^{(n-2)} \subset B^{(n-2)}$ and $A^{(n-2)}$ is left (resp. right) Arens irregular, then

$$A^{(n-2)} = \tilde{Z}_{B^{(n)}}^\ell(A^{(n)}) \text{ (resp. } A^{(n-2)} = \tilde{Z}_{B^{(n)}}^r(A^{(n)})).$$

Proof. (1) Assume that $(b_\alpha^{(n)})_\alpha \subseteq B^{(n)}$ such that $b_\alpha^{(n)} \xrightarrow{w^*} b^{(n)}$ in $B^{(n)}$. Let $b^{(n+1)} \in B^{(n+1)}$. Since $B^{(n+1)} = B^{(n-1)} \oplus B^\perp$, there are $b^{(n-1)} \in B^{(n-1)}$ and $t \in B^\perp$ such that $b^{(n+1)} = (b^{(n-1)}, t)$. Then for every $a^{(n-2)} \in A^{(n-2)}$, we have

$$\begin{aligned} \langle b^{(n+1)}, a^{(n-2)} b_\alpha^{(n)} \rangle &= \langle (b^{(n-1)}, t), a^{(n-2)} b_\alpha^{(n)} \rangle = \langle a^{(n-2)} b_\alpha^{(n)}, b^{(n-1)} \rangle \\ &\rightarrow \langle a^{(n-2)} b^{(n)}, b^{(n-1)} \rangle = \langle b^{(n+1)}, a^{(n-2)} b^{(n)} \rangle. \end{aligned}$$

It follows that $A^{(n-2)} \subset \tilde{Z}_{B^{(n)}}^\ell(A^{(n)})$.

(2) Proof is clear.

(3) Since $A^{(n-2)} \subset B^{(n-2)}$, it is clear that $\tilde{Z}_{B^{(n)}}^\ell(A^{(n)}) \subset Z_1(A^{(n)}) = A^{(n-2)}$.

Thus by part (1), since $\tilde{Z}_{B^{(n)}}^\ell(A^{(n)}) \subseteq Z_{B^{(n)}}^\ell(A^{(n)})$, we are done. $\square$

Example 3-7. Let G be a compact group. We know that $L^1(G) \subseteq M(G)$ and $L^1(G)$ is an ideal in $M(G)^{**}$. Since $L^1(G)$ is strongly Arens irregular, by preceding theorem, we conclude that

$$\tilde{Z}_{M(G)^{**}}^\ell(L^1(G)^{**}) \subseteq Z_{M(G)^{**}}^\ell(L^1(G)^{**}) \subseteq Z_1^\ell(L^1(G)^{**}) = L^1(G).$$

By Theorem 3-6, we have $L^1(G) \subseteq \tilde{Z}_{M(G)^{**}}^\ell(L^1(G)^{**})$. Thus we conclude that

$$L^1(G) = \tilde{Z}_{M(G)^{**}}^\ell(L^1(G)^{**}).$$

It is similar to above discussion that

$$L^1(G) = \tilde{Z}_{M(G)^{**}}^r(L^1(G)^{**}).$$

Theorem 3-8. Let B be a Banach A -bimodule. Then for every even number $n \geq 0$, we have the following assertions.

- (1) If $B^{(n+1)} B^{(n)} = A^{(n+1)}$ and $\tilde{Z}_{A^{(n)}}^\ell(B^{(n)}) = B^{(n)}$, then A is reflexive.
- (2) Let $e^{(n)} \in A^{(n)}$ be a left unit $A^{(n)}$ -module for $B^{(n)}$ and $e^{(n)} \in \tilde{Z}_{B^{(n)}}^\ell(A^{(n)})$. Then B is reflexive.

Proof. 1) Let $(a_\alpha^{(n)})_\alpha \subseteq A^{(n)}$ and $a_\alpha^{(n)} \xrightarrow{w^*} a^{(n)}$. Let $a^{(n+1)} \in A^{(n+1)}$. Since $B^{(n+1)} B^{(n)} \subseteq A^{(n+1)}$, there are $b^{(n+1)} \in B^{(n+1)}$ and $b^{(n)} \in B^{(n)}$ such that $a^{(n+1)} = b^{(n+1)} b^{(n)}$. Thus, we have

$$\begin{aligned} \langle a^{(n+1)}, a_\alpha^{(n)} \rangle &= \langle b^{(n+1)} b^{(n)}, a_\alpha^{(n)} \rangle = \langle b^{(n+1)}, b^{(n)} a_\alpha^{(n)} \rangle \rightarrow \langle b^{(n+1)}, b^{(n)} a^{(n)} \rangle \\ &= \langle a^{(n+1)}, a^{(n)} \rangle. \end{aligned}$$

We conclude that $A^{(n)}$ is reflexive, and so A is reflexive.

2) Let $(b_\alpha^{(n)})_\alpha \subseteq B^{(n)}$ and $b_\alpha^{(n)} \xrightarrow{w^*} b^{(n)}$. Let $b^{(n+1)} \in B^{(n+1)}$. Then

$$\langle b^{(n+1)}, b_\alpha^{(n)} \rangle = \langle b^{(n+1)}, e^{(n)} b_\alpha^{(n)} \rangle \rightarrow \langle b^{(n+1)}, e^{(n)} b^{(n)} \rangle = \langle b^{(n+1)}, b^{(n)} \rangle.$$

Thus B is reflexive. $\square$

Corollary 3-9. Assume that B is a Banach A – bimodule and B is reflexive. If $B^*B = A^*$, then A is reflexive.

Corollary 3-10. Assume that A is a Banach algebra and A^{**} has a left unit such that $\tilde{Z}_1^\ell(A^{**})$ consisting of it. Then A is reflexive.

For a Banach A – bimodule B and every members of A^{**} or B^{**} , we introduce locally weak topological centers of A^{**} and B^{**} and we establish some relations of them with reflexivity of A or B and Arens regularity of left and right module actions. Let B be a Banach left A – module. Suppose that $a'' \in A^{**}$ and $(a''_\alpha)_\alpha \subseteq A^{**}$ such that $a''_\alpha \xrightarrow{w^*} a''$. Then we say that $a'' \rightarrow \pi_\ell^{***}(a'', b'')$ is *weak* – to – weak** point continuous, if we have $\pi_\ell^{***}(a''_\alpha, b'') \xrightarrow{w^*} \pi_\ell^{***}(a'', b'')$. Suppose that B is a Banach A – bimodule. Assume that $a'' \in A^{**}$. Then we define the locally topological center of a'' on B^{**} as follows

$$Z_{a''}^\ell(B^{**}) = \{b'' \in B^{**} : a'' \rightarrow \pi_\ell^{***}(b'', a'') \text{ is weak}^* - \text{weak}^* \text{ point continuous}\},$$

$$Z_{a''}^r(B^{**}) = \{b'' \in B^{**} : b'' \rightarrow \pi_\ell^{t***}(a'', b'') \text{ is weak}^* - \text{weak}^* \text{ point continuous}\}.$$

The definition of $Z_{b''}^\ell(A^{**})$ and $Z_{b''}^r(A^{**})$ where $b'' \in B^{**}$ are similar. It is clear that

$$\begin{aligned} \bigcap_{a'' \in A^{**}} Z_{a''}^{\ell,r}(B^{**}) &= Z_{A^{**}}^{\ell,r}(B^{**}), \\ \bigcap_{b'' \in B^{**}} Z_{b''}^{\ell,r}(A^{**}) &= Z_{B^{**}}^{\ell,r}(A^{**}). \end{aligned}$$

Definition 3-11. Let B be a Banach A – bimodule. Assume that $a'' \in A^{**}$. Then we define the locally weak topological center of a'' on B^{**} in the following

$$\tilde{Z}_{a''}^\ell(B^{**}) = \{b'' \in B^{**} : a'' \rightarrow b''a'' \text{ is weak}^* - \text{weak point continuous}\},$$

$$\tilde{Z}_{a''}^r(B^{**}) = \{b'' \in B^{**} : a'' \rightarrow a''b'' \text{ is weak}^* - \text{weak point continuous}\}.$$

The definition of $\tilde{Z}_{b''}^\ell(A^{**})$ and $\tilde{Z}_{b''}^r(A^{**})$ where $b'' \in B^{**}$ are similar.

It is clear that

$$\begin{aligned} \bigcap_{a'' \in A^{**}} \tilde{Z}_{a''}^{\ell,r}(B^{**}) &= \tilde{Z}_{A^{**}}^{\ell,r}(B^{**}), \\ \bigcap_{b'' \in B^{**}} \tilde{Z}_{b''}^{\ell,r}(A^{**}) &= \tilde{Z}_{B^{**}}^{\ell,r}(A^{**}). \end{aligned}$$

Theorem 3-12. Let B be a Banach A – bimodule and let A^{**} has a right unit e'' such that $\tilde{Z}_{e''}^\ell(B^{**}) = B^{**}$. If B^* factors on the left with respect to A , then B is reflexive.

Proof. Let $(e_\alpha)_\alpha \subseteq A$ be a RBAI for A such that $e_\alpha \xrightarrow{w^*} e''$. Since B^* factors on the left with respect to A , for all $b' \in B^*$ there are $a \in A$ and $x' \in B^*$ such that $x'a = b'$. Then for all $b'' \in B^{**}$ we have

$$\begin{aligned} \langle \pi_\ell^{***}(e'', b''), b' \rangle &= \langle e'', \pi_\ell^{**}(b'', b') \rangle = \lim_\alpha \langle \pi_\ell^{**}(b'', b'), e_\alpha \rangle \\ &= \lim_\alpha \langle b'', \pi_\ell^*(b', e_\alpha) \rangle = \lim_\alpha \langle b'', \pi_\ell^*(x'a, e_\alpha) \rangle \\ &= \lim_\alpha \langle b'', \pi_\ell^*(x', ae_\alpha) \rangle = \lim_\alpha \langle \pi_\ell^{**}(b'', x'), ae_\alpha \rangle \\ &= \langle \pi_\ell^{**}(b'', x'), a \rangle = \langle b'', b' \rangle. \end{aligned}$$

Thus $\pi_\ell^{***}(e'', b'') = b''$ consequently B^{**} has left unit A^{**} – module. Since $\tilde{Z}_{e''}^\ell(B^{**}) = B^{**}$, it is clear that B is reflexive. $\square$

Assume that B is a Banach A – bimodule and $e'' \in A^{**}$ is a mixed unit for A^{**} . We say that e'' is left mixed unit for B^{**} , if $\pi_\ell^{***}(e'', b'') = \pi_\ell^{t***}(b'', e'') = b''$ for all $b'' \in B^{**}$.

The definition of the right mixed unit for B is similar. So, it is clear that $e^{(n)} \in A^{(n)}$ is a left (resp. right) mixed unit for $B^{(n)}$ if and only if $Z_{e^{(n)}}^\ell(B^{(n)}) = B^{(n)}$ (resp. $Z_{e^{(n)}}^r(B^{(n)}) = B^{(n)}$) whenever n is even.

Lemma 3-13. Let B be a Banach A – bimodule. Then for every even number $n \geq 2$, we have

- (1) If $B^{(n-2)}$ has a bounded right approximate identity (= BRAI) as $A^{(n-2)}$ – module, then $B^{(n)}$ has right unit as $A^{(n)}$ – module.
- (2) If $B^{(n-1)}$ has a bounded BRAI as $A^{(n-1)}$ – module, then $B^{(n)}$ has left unit as $A^{(n+1)}$ – module.

Proof. 1) Let $(e_\alpha^{(n-2)})_\alpha \subseteq A^{(n-2)}$ be a BRAI for $B^{(n-2)}$. By Goldstines theorem there is a right unit $e^{(n)} \in A^{(n)}$ such that it is *weak** closure of $(e_\alpha^{(n-2)})_\alpha$. Without lose generality, let $e_\alpha^{(n-2)} \xrightarrow{w^*} e^{(n)}$ in $A^{(n)}$. Assume that $b^{(n-1)} \in B^{(n-1)}$ and $b^{(n-2)} \in B^{(n-2)}$. Then

$$\begin{aligned} \langle b^{(n-1)}, b^{(n-2)} \rangle &= \lim_\alpha \langle b^{(n-1)}, b^{(n-2)} e_\alpha^{(n-2)} \rangle = \lim_\alpha \langle e_\alpha^{(n-2)}, b^{(n-1)} b^{(n-2)} \rangle \\ &= \langle e^{(n)}, b^{(n-1)} b^{(n-2)} \rangle. \end{aligned}$$

Then $e^{(n)} b^{(n-1)} = b^{(n-1)}$. Now let $b^{(n)} \in B^{(n)}$. Then we have

$$\langle b^{(n)} e^{(n)}, b^{(n-1)} \rangle = \langle b^{(n)}, e^{(n)} b^{(n-1)} \rangle = \langle b^{(n)}, b^{(n-1)} \rangle.$$

We conclude that $b^{(n)} e^{(n)} = b^{(n)}$.

2) Proof is similar to (1). $\square$

Corollary 3-14. Let B be a Banach A – bimodule and suppose that $n \geq 2$ is even number. Then $B^{(n-2)}$ has a BAI $A^{(n-2)}$ – module if and only if $B^{(n)}$ has a unit $A^{(n)}$ – module.

Theorem 3-15. Let B be a Banach A – bimodule. Then for every even number $n \geq 2$, we have the following assertions.

- (1) If $e^{(n)} \in A^{(n)}$ is a right unit for $B^{(n)}$ and $Z_{e^{(n)}}^\ell(B^{(n)}) = B^{(n)}$, then
$$A^{(n-2)}B^{(n-1)} = B^{(n-1)}.$$
- (2) If $e^{(n)} \in A^{(n)}$ is a left unit for $B^{(n)}$ and $Z_{e^{(n)}}^r(B^{(n)}) = B^{(n)}$, then
$$B^{(n-1)}A^{(n-2)} = B^{(n-1)}.$$
- (3) Suppose that $e^{(n)} \in A^{(n)}$ is a right unit for $B^{(n)}$ and $\tilde{Z}_{e^{(n)}}^\ell(B^{(n)}) = B^{(n)}$. Then $A^{(n-2)}B^{(n-1)} = B^{(n-1)}$ and *weak** closure of $A^{(n-2)}B^{(n+1)}$ is $B^{(n+1)}$.
- (4) Suppose that $e^{(n)} \in A^{(n)}$ is a left unit for $B^{(n)}$ and $\tilde{Z}_{e^{(n)}}^r(B^{(n)}) = B^{(n)}$. Then $B^{(n-1)}A^{(n-2)} = B^{(n-1)}$ and *weak** closure of $B^{(n+1)}A^{(n-2)}$ is $B^{(n+1)}$.

Proof. 1) Since $e^{(n)}$ is a right unit for $B^{(n)}$, by using Lemma 3-13, there is a BRAI as $(e_\alpha^{(n-2)})_\alpha \subseteq A^{(n-2)}$ for $A^{(n-2)}$ such that $e_\alpha^{(n-2)} \xrightarrow{w^*} e^{(n)}$ in $A^{(n)}$. Let $b^{(n)} \in B^{(n)}$. Since $Z_{e^{(n)}}^\ell(B^{(n)}) = B^{(n)}$, $b^{(n)}e_\alpha^{(n-2)} \xrightarrow{w^*} b^{(n)}e^{(n)} = b^{(n)}$. Then for all $b^{(n-1)} \in B^{(n-1)}$, we have

$$\begin{aligned} \langle b^{(n)}, e_\alpha^{(n-2)}b^{(n-1)} \rangle &= \langle b^{(n)}e_\alpha^{(n-2)}, b^{(n-1)} \rangle \rightarrow \langle b^{(n)}e^{(n)}, b^{(n-1)} \rangle = \\ &= \langle b^{(n)}, b^{(n-1)} \rangle. \end{aligned}$$

It follows that $e_\alpha^{(n-2)}b^{(n-1)} \xrightarrow{w} b^{(n-1)}$. Consequently, by using Cohen factorization theorem, we have $A^{(n-2)}B^{(n-1)} = B^{(n-1)}$.

2) Proof is similar to (1).

3) Since $\tilde{Z}_{e^{(n)}}^\ell(B^{(n)}) \subseteq Z_{e^{(n)}}^\ell(B^{(n)})$, it is clear that $A^{(n-2)}B^{(n-1)} = B^{(n-1)}$. We prove that the *weak** closure of $A^{(n-2)}B^{(n+1)}$ is $B^{(n+1)}$. Since $e^{(n)}$ is a right unit for $B^{(n)}$, by using Lemma 3-13, there is $(e_\alpha^{(n-2)})_\alpha \subseteq A^{(n-2)}$ such that $e_\alpha^{(n-2)} \xrightarrow{w^*} e^{(n)}$ in $A^{(n)}$. Since $\tilde{Z}_{e^{(n)}}^r(B^{(n)}) = B^{(n)}$, $b^{(n)}e_\alpha^{(n-2)} \xrightarrow{w} b^{(n)}e^{(n)} = b^{(n)}$ for all $b^{(n)} \in B^{(n)}$. Then for every $b^{(n+1)} \in B^{(n+1)}$, we have

$$\langle e_\alpha^{(n-2)}b^{(n+1)}, b^{(n)} \rangle = \langle b^{(n+1)}, b^{(n)}e_\alpha^{(n-2)} \rangle \rightarrow \langle b^{(n+1)}, b^{(n)}e^{(n)} \rangle = \langle b^{(n+1)}, b^{(n)} \rangle.$$

It follow that $e_\alpha^{(n-2)}b^{(n+1)} \xrightarrow{w^*} b^{(n+1)}$, and so proof hold.

4) Proof is similar to (3). □

Corollary 3-16. Assume that A is a Banach algebra. Then we have the following assertions.

- (1) If $Z_{e''}^\ell(A^{**}) = A^{**}$, then A^* factors on the right where e'' is a right unite for A^{**} .

(2) If $Z_{e''}^r(A^{**}) = A^{**}$, then A^* factors on the left where e'' is a left unite for A^{**} .

Corollary 3-17. Assume that A is a Banach algebra and has a BAI . If A^* not factors on the one side, then A is not Arens regular.

Proof. Since $Z_1^\ell(A^{**}) \subseteq Z_{e''}^\ell(A^{**})$, by Theorem 3-15, proof hold. $\square$

Example 3-18. Let G be a locally compact group and e'' be a mixed unit for $L^1(G)^{**}$. Then we have $Z_{e''}^\ell(L^1(G)^{**}) \neq L^1(G)^{**}$ and $Z_{e''}^r(L^1(G)^{**}) \neq L^1(G)^{**}$.

Theorem 3-19. Assume that A is a Banach algebra. Then $n \geq 0$, we have the following assertions.

- (1) If $A^{(n)}$ has a BAI , then $\tilde{Z}_{A^{(n+2)}}^\ell(A^{(n+3)}) \subseteq A^{(n+1)}$. Moreover, for every odd number n , if $A^{(n+3)}A^{(n+1)} \subseteq A^{(n+1)}$, then $\tilde{Z}_{A^{(n+2)}}^\ell(A^{(n+3)}) = A^{(n+1)}$.
- (2) If $A^{(n+2)}$ has a left unit, then $\tilde{Z}_{A^{(n+3)}}^r(A^{(n+2)}) \subseteq A^{(n)}$. Moreover, for every even number n , if $A^{(n+4)}A^{(n)} \subseteq A^{(n+1)}$, then $\tilde{Z}_{A^{(n+3)}}^r(A^{(n+2)}) = A^{(n)}$.

Proof. 1) Assume that $A^{(n)}$ has a BAI as $(e_\alpha^{(n)})_\alpha \subseteq A^{(n)}$ such that $e_\alpha^{(n)} \xrightarrow{w^*} e^{(n+2)}$ in $A^{(n+2)}$ where $e^{(n+2)}$ is a right unite in $A^{(n+2)}$. Let $a^{(n+3)} \in \tilde{Z}_{A^{(n+2)}}^\ell(A^{(n+3)})$ and $(a_\alpha^{(n+2)})_\alpha \subseteq A^{(n+2)}$ such that $a_\alpha^{(n+2)} \xrightarrow{w^*} a^{(n+2)}$ in $A^{(n+2)}$. Then we have

$$\begin{aligned} \langle a^{(n+3)}, a_\alpha^{(n+2)} \rangle &= \langle a^{(n+3)}, a_\alpha^{(n+2)} e^{(n+2)} \rangle = \langle a^{(n+3)} a_\alpha^{(n+2)}, e^{(n+2)} \rangle \\ &= \langle e^{(n+2)}, a^{(n+3)} a_\alpha^{(n+2)} \rangle \rightarrow \langle e^{(n+2)}, a^{(n+3)} a^{(n+2)} \rangle = \langle a^{(n+3)}, a^{(n+2)} \rangle. \end{aligned}$$

It follows that $a^{(n+3)} : A^{(n+3)} \rightarrow \mathbb{C}$ is *weak** continuous, and so $a^{(n+3)} \in A^{(n+1)}$. Consequently, we have $\tilde{Z}_{A^{(n+2)}}^\ell(A^{(n+3)}) \subseteq A^{(n+1)}$.

Now let $A^{(n+3)}A^{(n+1)} \subseteq A^{(n+1)}$. We show that the mapping $a^{(n+2)} \rightarrow a^{(n+1)}a^{(n+2)}$ is *weak*-to-weak* continuous for all $a^{(n+1)} \in A^{(n+1)}$. Assume that $(a_\alpha^{(n+2)})_\alpha \subseteq A^{(n+2)}$ such that $a_\alpha^{(n+2)} \xrightarrow{w^*} a^{(n+2)}$ in $A^{(n+2)}$. Let $a^{(n+3)} \in A^{(n+3)}$. Then we have

$$\begin{aligned} \langle a^{(n+3)}, a^{(n+1)} a_\alpha^{(n+2)} \rangle &= \langle a^{(n+3)} a^{(n+1)}, a_\alpha^{(n+2)} \rangle = \langle a_\alpha^{(n+2)}, a^{(n+3)} a^{(n+1)} \rangle \\ &\rightarrow \langle a^{(n+2)}, a^{(n+3)} a^{(n+1)} \rangle = \langle a^{(n+3)}, a^{(n+1)} a^{(n+2)} \rangle. \end{aligned}$$

It follows that $a^{(n+1)} \in \tilde{Z}_{A^{(n+2)}}^\ell(A^{(n+3)})$, and so $\tilde{Z}_{A^{(n+2)}}^\ell(A^{(n+3)}) = A^{(n+1)}$.

2) It is similar to (1). $\square$

Corollary 3-20. Assume that A is a Banach algebra. Then

- (1) If A has a BAI and $\tilde{Z}_{A^{**}}^\ell(A^{***}) = A^{***}$. Then A is reflexive.
- (2) If A has a left unit and $\tilde{Z}_{A^{***}}^r(A^{**}) = A^{**}$. Then A is reflexive.

Example 3-21. Let G be a locally compact infinite group. Then we have the following statements.

- (1) $\tilde{Z}^\ell_{L^1(G)**}(L^1(G)^{***}) \subseteq L^1(G)^*$ and $\tilde{Z}^r_{L^1(G)^{***}}(L^1(G)^{**}) \subseteq L^1(G)$ where multiplications are with respect to the first and second Arens products, respectively.
- (2) Since $c_0^* = \ell^1$, we have $\tilde{Z}^\ell_{\ell^\infty}((\ell^\infty)^*) \subseteq \ell^\infty$ and $\tilde{Z}^r_{(\ell^\infty)^*}(\ell^\infty) \subseteq c_0$ where multiplications in c_0^{**} and ℓ^∞ coincide.

Theorem 3-22. Assume that B is a Banach A –*bimodule*. Then for all $n \geq 1$, we have

- i) If $B^{(n)}B^{(n-1)} = A^{(n-1)}$, then $Z^\ell_{B^{(n)}}(A^{(n)}) \subseteq Z_1(A^{(n)})$.
- ii) If $B^{(n+1)}B^{(n)} = A^{(n+1)}$, then $Z^r_{B^{(n+1)}}(A^{(n)}) \subseteq Z_1(A^{(n)})$.

Proof. i) Let $a^{(n)} \in Z_{B^{(n)}}(A^{(n)})$. We show that $x^{(n)} \rightarrow a^{(n)}x^{(n)}$ from $A^{(n)}$ into $A^{(n)}$ is *weak*–to–weak** continuous. Suppose that $(x_\alpha^{(n)})_\alpha \subseteq A^{(n)}$ such that $x_\alpha^{(n)} \xrightarrow{w^*} x^{(n)}$ in $A^{(n)}$. First we show that $x_\alpha^{(n)}b^{(n)} \xrightarrow{w^*} x^{(n)}b^{(n)}$ in $B^{(n)}$. Let $b^{(n-1)} \in B^{(n-1)}$. Then

$$\begin{aligned} \langle x_\alpha^{(n)}b^{(n)}, b^{(n-1)} \rangle &= \langle x_\alpha^{(n)}, b^{(n)}b^{(n-1)} \rangle \rightarrow \langle x^{(n)}, b^{(n)}b^{(n-1)} \rangle \\ &= \langle x^{(n)}b^{(n)}, b^{(n-1)} \rangle. \end{aligned}$$

Now let $a^{(n-1)} \in A^{(n-1)}$. Then we have

$$\begin{aligned} \langle a^{(n)}x_\alpha^{(n)}, a^{(n-1)} \rangle &= \langle a^{(n)}x_\alpha^{(n)}, b^{(n)}b^{(n-1)} \rangle = \langle a^{(n)}x_\alpha^{(n)}b^{(n)}, b^{(n-1)} \rangle \\ &\rightarrow \langle a^{(n)}x^{(n)}b^{(n)}, b^{(n-1)} \rangle = \langle a^{(n)}x^{(n)}, b^{(n)}b^{(n-1)} \rangle \\ &= \langle a^{(n)}x^{(n)}, a^{(n-1)} \rangle. \end{aligned}$$

It follows that $a^{(n)} \in Z_1(A^{(n)})$.

ii) Proof is similar to (1). □

Corollary 3-23. Assume that B is a Banach A –*bimodule* with left and right module actions π_ℓ and π_r , respectively. Then

- (1) If π_ℓ^{**} is surjective, then $Z^\ell_{B^{**}}(A^{**}) \subseteq Z_1(A^{**})$
- (2) If π_r^{**} is surjective, then $Z^r_{B^{**}}(A^{**}) \subseteq Z_2(A^{**})$

Example 3-24. Let G be a locally compact group and $n \geq 3$. Then

- (1) $Z^\ell_{L^1(G)^{(n)}}(L^1(G)^{**}) = L^1(G)$, see [8].
- (2) $Z^\ell_{M(G)^{(n)}}(L^1(G)^{**}) = L^1(G)$.
- (3) $Z^\ell_{M(G)^{(n)}}(M(G)^{**}) = M(G)$.

Theorem 3-25. Let B be a Banach A –*bimodule*. Then for every even number $n \geq 2$, we have the following assertions.

- (1) If $B^{(n-1)}$ has a $BLAI$ as $A^{(n-2)}$ - module such that $weak^*$ convergence to $e^{(n)} \in A^{(n)}$ and $\tilde{Z}_{e^{(n)}}^r(B^{(n-1)}) = B^{(n-1)}$, then $B^{(n)}$ has right unit as $A^{(n)}$ - module.
- (2) If $B^{(n-1)}$ has a $BRAI$ as $A^{(n-1)}$ - module such that $weak^*$ convergence to $e^{(n+1)} \in A^{(n+1)}$ and $\tilde{Z}_{e^{(n+1)}}^\ell(B^{(n-1)}) = B^{(n-1)}$, then $B^{(n)}$ has left unit as $A^{(n+1)}$ - module.
- (3) If $B^{(n-2)}$ has a $BLAI$ as $A^{(n-2)}$ - module such that $weak^*$ convergence to $e^{(n)} \in A^{(n)}$ and $\tilde{Z}_{e^{(n)}}^\ell(B^{(n-1)}) = B^{(n-1)}$, then $B^{(n)}$ has left unit as $A^{(n)}$ - module.
- (4) Suppose that $B^{(n-1)}$ has a $BRAI$ as $A^{(n-1)}$ - module such that $weak^*$ convergence to $e^{(n+1)} \in A^{(n+1)}$. Let $\tilde{Z}_{e^{(n+1)}}^\ell(B^{(n-1)}) = B^{(n-1)}$. Then $e^{(n+1)}$ is a left unit as $A^{(n+1)}$ - module for $B^{(n)}$.

Proof. 1) Suppose that $(e_\alpha^{(n-2)})_\alpha \subseteq A^{(n-2)}$ is a $BLAI$ for $B^{(n-1)}$. Then there is a right unit element $e^{(n)} \in A^{(n)}$ for $A^{(n)}$ such that $e_\alpha^{(n-2)} \xrightarrow{w^*} e^{(n)}$ in $A^{(n)}$. Let $b^{(n-1)} \in B^{(n-1)}$. Since $\tilde{Z}_{e^{(n)}}^r(B^{(n-1)}) = B^{(n-1)}$, we have $e_\alpha^{(n-2)} b^{(n-1)} \xrightarrow{w} e^{(n)} b^{(n-1)}$. Then for every $b^{(n)} \in B^{(n)}$, we have

$$\begin{aligned} \langle b^{(n)} e^{(n)}, b^{(n-1)} \rangle &= \langle b^{(n)}, e^{(n)} b^{(n-1)} \rangle = \lim_\alpha \langle b^{(n)}, e_\alpha^{(n-2)} b^{(n-1)} \rangle \\ &= \langle b^{(n)}, b^{(n-1)} \rangle. \end{aligned}$$

The proof of (2), (3) and (4) is similar to (1). $\square$

Corollary 3-26. Let B be a Banach A - bimodule. Then for every even number $n \geq 2$, we have the following assertions.

- (1) If $B^{(n-1)}$ has a BAI as $A^{(n-2)}$ - module such that $weak^*$ convergence to $e^{(n)} \in A^{(n)}$ and $\tilde{Z}_{e^{(n)}}^r(B^{(n-1)}) = B^{(n-1)}$, then $B^{(n)}$ is unital as $A^{(n)}$ - module.
- (2) If $B^{(n-1)}$ has a BAI as $A^{(n-1)}$ - module such that $weak^*$ convergence to $e^{(n+1)} \in A^{(n+1)}$ and $\tilde{Z}_{e^{(n+1)}}^\ell(B^{(n-1)}) = B^{(n-1)}$, then $B^{(n)}$ is unital as $A^{(n+1)}$ - module.
- (3) If $B^{(n-2)}$ has a BAI as $A^{(n-2)}$ - module such that $weak^*$ convergence to $e^{(n)} \in A^{(n)}$ and $\tilde{Z}_{e^{(n)}}^\ell(B^{(n-1)}) = B^{(n-1)}$, then $B^{(n)}$ is unital as $A^{(n)}$ - module.

Proof. By using Lemma 3-13 and Theorem 3-25, proof is hold. $\square$

Theorem 3-27. Let A be a Banach algebra and $n > 0$ be even number.

- i) Assume that $A^{(n)}$ has a right (resp. left) unit. Then $Z_{A^{(n)}}^\ell(A^{(n+1)}) = A^{(n-1)}$ (resp. $Z_{A^{(n)}}^r(A^{(n+1)}) = A^{(n-1)}$) where multiplication is to the first (resp. second) Arens product.
- ii) Assume that $A^{(n-1)}$ has a right unit as $A^{(n-2)}$ -module. Then $Z_{A^{(n-1)}}^r(A^{(n)}) = A^{(n-2)}$ where multiplication is to the first Arens product.

Proof. Assume that $(a_\alpha^{(n)})_\alpha \subseteq A^{(n)}$ such that $a_\alpha^{(n)} \xrightarrow{w^*} a^{(n)}$ in $A^{(n)}$. Let $a^{(n+1)} \in Z_{A^{(n)}}^\ell(A^{(n+1)})$ and $e^{(n)} \in A^{(n)}$ be a left unit for $A^{(n)}$. Then

$$\begin{aligned} \langle a^{(n+1)}, a_\alpha^{(n)} \rangle &= \langle a^{(n+1)}, a_\alpha^{(n)} e^{(n)} \rangle = \langle a^{(n+1)} a_\alpha^{(n)}, e^{(n)} \rangle \rightarrow \langle a^{(n+1)} a^{(n)}, e^{(n)} \rangle \\ &= \langle a^{(n+1)}, a^{(n)} \rangle. \end{aligned}$$

It follows that $a^{(n+1)} \in (A^{(n+1)}, weak^*)^* = A^{(n-1)}$.

Next part is similar.

ii) Proof is clear. $\square$

Corollary 3-28. Let A be a Banach algebra with BAI Then $Z_{A^{**}}^\ell(A^{***}) = A^*$ (resp. $Z_{A^{**}}^r(A^{***}) = A^*$) where multiplication is to the first (resp. second) Arens product.

Example 3-29. Let G be a locally compact groups. Then

- i) $Z_{L^1(G)^{**}}^\ell(L^1(G)^{***}) = L^1(G)^*$ (resp. $Z_{L^1(G)^{**}}^r(L^1(G)^{***}) = L^1(G)^*$) where multiplication is to the first (resp. second) Arens product.
- ii) Let $n > 0$ be even number. Then $Z_{M(G)^{(n)}}^\ell(M(G)^{(n+1)}) = M(G)^{(n-1)}$ (resp. $Z_{M(G)^{(n)}}^r(M(G)^{(n+1)}) = M(G)^{(n-1)}$) where multiplication is to the first (resp. second) Arens product.

4. Weak amenability of Banach algebras

In this section, for even number $n \geq 2$, we show that if the weak topological center of $A^{(n)}$ with respect to the first Arens product be itself, then $A^{(n-2)}$ is weakly amenable whenever $A^{(n)}$ is weakly amenable. As corollary we show that if $\widetilde{wap}_\ell(A^{(n)}) \subseteq A^{(n+1)}$, then the weakly amenability of $A^{(n)}$ implies the weakly amenability of $A^{(n-2)}$ for each $n \geq 2$. For a Banach A – bimodule B , we deal with the question of when the second adjoint $D'' : A^{**} \rightarrow B^{***}$ of $D : A \rightarrow B^*$ is again a derivation. This problem has been studied in [5, 9, 15] and we try to give answer to this question, that is, if the left weak topological center of a Banach algebra A^{**} is A , then $D : A \rightarrow B^*$ is a derivation whenever $D'' : A^{**} \rightarrow B^{***}$ is a derivation.

At the beginning, we introduce some elementary notions and definitions as follows: Let B a Banach A – bimodule. A derivation from A into B is a bounded linear mapping $D : A \rightarrow B$ such that

$$D(xy) = xD(y) + D(x)y \text{ for all } x, y \in A.$$

The space of continuous derivations from A into B is denoted by $Z^1(A, B)$.

Easy example of derivations are the inner derivations, which are given for each $b \in B$ by

$$\delta_b(a) = ab - ba \text{ for all } a \in A.$$

The space of inner derivations from A into B is denoted by $N^1(A, B)$. A Banach algebra A is said to be a weakly amenable, when for every Banach A -bimodule B , the inner derivations are only derivations existing from A into B^* . It is clear that A is amenable if and only if $H^1(A, B^*) = Z^1(A, B^*)/N^1(A, B^*) = \{0\}$.

A Banach algebra A is said to be a amenable, if every derivation from A into A^* is inner. Similarly, A is weakly amenable if and only if

$$H^1(A, A^*) = Z^1(A, A^*)/N^1(A, A^*) = \{0\}.$$

In every parts of this section, $n \geq 0$ is an even number.

Theorem 4-1. Assume that A is a Banach algebra and $\tilde{Z}_1^\ell(A^{(n)}) = A^{(n)}$ where $n \geq 2$. If $A^{(n)}$ is weakly amenable, then $A^{(n-2)}$ is weakly amenable.

Proof. Suppose that $D \in Z^1(A^{(n-2)}, A^{(n-1)})$. First we show that

$$D'' \in Z^1(A^{(n)}, A^{(n+1)}).$$

Let $a^{(n)}, b^{(n)} \in A^{(n)}$ and let $(a_\alpha^{(n-2)})_\alpha, (b_\beta^{(n-2)})_\beta \subseteq A^{(n-2)}$ such that $a_\alpha^{(n-2)} \xrightarrow{w^*} a^{(n)}$ and $b_\beta^{(n-2)} \xrightarrow{w^*} b^{(n)}$, respectively. Since $\tilde{Z}_1^\ell(A^{(n)}) = A^{(n)}$ we have

$$\lim_{\alpha} \lim_{\beta} a_\alpha^{(n-2)} D(b_\beta^{(n-2)}) = a^{(n)} D''(b^{(n)}).$$

It is also clear that

$$\lim_{\alpha} \lim_{\beta} D(a_\alpha^{(n-2)}) b_\beta^{(n-2)} = D''(a^{(n)}) b^{(n)}.$$

Since D is continuous, we conclude that

$$D''(a^{(n)} b^{(n)}) = \lim_{\alpha} \lim_{\beta} D(a_\alpha^{(n-2)} b_\beta^{(n-2)}) = \lim_{\alpha} \lim_{\beta} a_\alpha^{(n-2)} D(b_\beta^{(n-2)}) +$$

$$\lim_{\alpha} \lim_{\beta} D(a_\alpha^{(n-2)}) b_\beta^{(n-2)} = a^{(n)} D''(b^{(n)}) + D''(a^{(n)}) b^{(n)}.$$

In above, we take limit in *weak** topology. Since $A^{(n)}$ is weakly amenable, D'' is inner. It follows that $D''(a^{(n)}) = a^{(n)} a^{(n+1)} - a^{(n+1)} a^{(n)}$ for some $a^{(n+1)} \in A^{(n+1)}$. Take $a^{(n-1)} = a^{(n+1)}|_{A^{(n-2)}}$ and $a^{(n-2)} \in A^{(n-2)}$. Then

$$D(a^{(n-2)}) = D''(a^{(n-2)}) = a^{(n-2)} a^{(n-1)} - a^{(n-1)} a^{(n-2)} = \delta_{a^{(n-1)}}(a^{(n-2)}).$$

Consequently, we have $H^1(A^{(n-2)}, A^{(n-1)}) = 0$, and so $A^{(n-2)}$ is weakly amenable. $\square$

Corollary 4-2. Let A be a Banach algebra and let $\widetilde{wap}_\ell(A^{(n-1)}) \subseteq A^{(n)}$ whenever $n \geq 1$. If $A^{(n)}$ is weakly amenable, then $A^{(n-2)}$ is weakly amenable.

Proof. Since $\widetilde{wap}_\ell(A^{(n-1)}) \subseteq A^{(n)}$, $\tilde{Z}_1^\ell(A^{(n)}) = A^{(n)}$. Then, by using Theorem 4-1, proof hold. $\square$

Corollary 4-3. Let A be a Banach algebra and $Z^\ell_{A^{(n)}}(A^{(n+1)}) = A^{(n+1)}$ where $n \geq 2$. If $A^{(n)}$ is weakly amenable, then $A^{(n-2)}$ is weakly amenable.

Corollary 4-4. Let A be a Banach algebra and let $D : A^{(n-2)} \rightarrow A^{(n-1)}$ be a derivation. Then $D'' : A^{(n)} \rightarrow A^{(n+1)}$ is a derivation, if $\tilde{Z}_1^\ell(A^{(n)}) = A^{(n)}$ for every $n \geq 2$.

Theorem 4-5. Let A be a Banach algebra and let B be a closed subalgebra of $A^{(n)}$ that is consisting of $A^{(n-2)}$. If $\tilde{Z}_1^\ell(B) = B$ and B is weakly amenable, then $A^{(n-2)}$ is weakly amenable.

Proof. Suppose that $D : A^{(n-2)} \rightarrow A^{(n-1)}$ is a derivation and $p : A^{(n+1)} \rightarrow B^*$ is the restriction map, defined by $P(a^{(n+1)}) = a^{(n+1)}|_B$ for every $a^{(n+1)} \in A^{(n+1)}$. Since $\tilde{Z}_1^\ell(B) = B$, $\bar{D} = P \circ D''|_B : B \rightarrow B'$ is a derivation. Since B is weakly amenable, there is $b' \in B^*$ such that $\bar{D} = \delta_{b'}$. We take $a^{(n-1)} = b'|_{A^{(n-2)}}$, then $D = \bar{D}$ on $A^{(n-2)}$. Consequently, we have $D = \delta_{a^{(n-1)}}$. $\square$

Corollary 4-6. Let A be a Banach algebra. If $A^{***}A^{**} \subseteq A^*$ and A is weakly amenable, then A^{**} is weakly amenable.

Proof. By using Corollary 2-4 and Theorem 3-2, proof hold. $\square$

Corollary 4-7. Let A be a Banach algebra and let $\tilde{Z}_1^\ell(A^{(n)})$ be weakly amenable whenever $n \geq 2$. Then $A^{(n-2)}$ is weakly amenable.

Theorem 4-8. Let B be a Banach A -bimodule and $D : A^{(n)} \rightarrow B^{(n+1)}$ be a derivation where $n \geq 0$. If $\tilde{Z}^\ell_{A^{(n+2)}}(B^{(n+2)}) = B^{(n+2)}$, then $D'' : A^{(n+2)} \rightarrow B^{(n+3)}$ is a derivation.

Proof. Let $x^{(n+2)}, y^{(n+2)} \in A^{(n+2)}$ and let $(x_\alpha^{(n)})_\alpha, (y_\beta^{(n)})_\beta \subseteq A^{(n)}$ such that $x_\alpha^{(n)} \xrightarrow{w^*} x^{(n+2)}$ and $y_\beta^{(n)} \xrightarrow{w^*} y^{(n+2)}$ in $A^{(n+2)}$. Then for all $b^{(n+2)} \in B^{(n+2)}$, we have $b^{(n+2)}x_\alpha^{(n)} \xrightarrow{w} b^{(n+2)}x^{(n+2)}$. Consequently, since $\tilde{Z}^\ell_{A^{(n+2)}}(B^{(n+2)}) = B^{(n+2)}$, we have

$$\begin{aligned} \langle x_\alpha^{(n)} D''(y^{(n+2)}), b^{(n+2)} \rangle &= \langle D''(y^{(n+2)}), b^{(n+2)} x_\alpha^{(n)} \rangle \\ &\rightarrow \langle D''(y^{(n+2)}), b^{(n+2)} x^{(n+2)} \rangle \\ &= \langle x^{(n+2)} D''(y^{(n+2)}), b^{(n+2)} \rangle. \end{aligned}$$

Also we have the following equality

$$\begin{aligned} \langle D''(x^{(n+2)}) y_\beta^{(n)}, b^{(n+2)} \rangle &= \langle D''(x^{(n+2)}), y_\beta^{(n)} b^{(n+2)} \rangle \\ &\rightarrow \langle D''(x^{(n+2)}), y^{(n+2)} b^{(n+2)} \rangle \\ &= \langle D''(x^{(n+2)}) y^{(n+2)}, b^{(n+2)} \rangle. \end{aligned}$$

Since D is continuous, it follows that

$$\begin{aligned} D''(x^{(n+2)}y^{(n+2)}) &= \lim_{\alpha} \lim_{\beta} D(x_{\alpha}^{(n)}y_{\beta}^{(n)}) = \lim_{\alpha} \lim_{\beta} x_{\alpha}^{(n)}D(y_{\beta}^{(n)}) + \\ \lim_{\alpha} \lim_{\beta} D(x_{\alpha}^{(n)}y_{\beta}^{(n)}) &= x^{(n+2)}D''(y^{(n+2)}) + D''(x^{(n+2)})y^{(n+2)} \end{aligned}$$

□

Corollary 4-9. Let B be a Banach A -bimodule and $\tilde{Z}_{A^{**}}^{\ell}(B^{**}) = B^{**}$. Then, if $H^1(A, B^*) = 0$, it follows that $H^1(A^{**}, B^{***}) = 0$.

Corollary 4-10. Let B be a Banach A -bimodule and $\tilde{Z}_{A^{**}}^{\ell}(B^{**}) = B^{**}$. Let $D : A \rightarrow B^*$ be a derivation. Then $D''(A^{**})B^{**} \subseteq A^*$.

Proof. By using Theorem 4-8 and [15, Corollary 4-3], proof is hold.

□

Corollary 4-11. Let B be a Banach A -bimodule and let $D : A \rightarrow B^*$ be a derivation such that D is surjective. Then $Z_{A^{**}}^{\ell}(B^{**}) = \tilde{Z}_{A^{**}}^{\ell}(B^{**})$.

Proof. By using Theorem 3-2 and [15, Corollary 4-3], proof is hold.

□

Problems:

- Let G be a locally compact infinite group. Find the sets $\tilde{Z}_1^{\ell}(L^1(G)^{**}) = ?$ and $\tilde{Z}_1^{\ell}(M(G)^{**}) = ?$
- Assume that $n > 0$ is even number and A and B are Banach algebra. If $\tilde{Z}^{\ell}(A^{(n)}) = A^{(n)}$ and $\tilde{Z}^{\ell}(B^{(n)}) = B^{(n)}$, what can say about $\tilde{Z}^{\ell}(A^{(n)} \otimes B^{(n)})$?
- Assume that A and B are Banach algebras. If $\tilde{Z}^{\ell}(A^{(n)} \otimes B^{(n)}) = A^{(n)} \otimes B^{(n)}$, what can say about $\tilde{Z}^{\ell}(A^{(n)})$ and $\tilde{Z}^{\ell}(B^{(n)})$?
- Let G be a locally compact group and $n \geq 3$. By notice to Theorem 3-20 and Example 3-22, what can say about to the following sets.

$$\tilde{Z}_{L^1(G)^{(n)}}^{\ell}(L^1(G)^{**}), \quad \tilde{Z}_{M(G)^{(n)}}^{\ell}(L^1(G)^{**}), \quad \tilde{Z}_{M(G)^{(n)}}^{\ell}(M(G)^{**}).$$

REFERENCES

1. R. E. Arens, *The adjoint of a bilinear operation*, Proc. Amer. Math. Soc. **2** (1951), 839-848.
2. N. Arian, *A simple condition ensuring the Arens regularity of bilinear mappings*, Proc. Amer. Math. Soc. **84** (4) (1982), 525-532.
3. J. Baker, A.T. Lau, J.S. Pym *Module homomorphism and topological centers associated with weakly sequentially compact Banach algebras*, Journal of Functional Analysis. **158** (1998), 186-208.
4. F. F. Bonsall, J. Duncan, *Complete normed algebras*, Springer-Verlag, Berlin 1973.
5. H. G. Dales, A. Rodrigues-Palacios, M.V. Velasco, *The second transpose of a derivation*, J. London. Math. Soc. **2** 64 (2001) 707-721.
6. H. G. Dales, *Banach algebra and automatic continuity*, Oxford 2000.
7. N. Dunford, J. T. Schwartz, *Linear operators. I*, Wiley, New york 1958.
8. M. Eshaghi Gordji, M. Filali, *Arens regularity of module actions*, Studia Math. **181** 3 (2007), 237-254.

9. M. Eshaghi Gordji, M. Filali, *Weak amenability of the second dual of a Banach algebra*, Studia Math. **182** 3 (2007), 205-213.
10. K. Haghnejad Azar, A. Riazi, *Arens regularity of bilinear forms and unital Banach module space*, (Submitted).
11. E. Hewitt, K. A. Ross, *Abstract harmonic analysis*, Springer, Berlin, Vol I 1963.
12. E. Hewitt, K. A. Ross, *Abstract harmonic analysis*, Springer, Berlin, Vol II 1970.
13. A. T. Lau, V. Losert, *On the second Conjugate Algebra of locally compact groups*, J. London Math. Soc. **37** (2)(1988), 464-480.
14. A. T. Lau, A. Ülger, *Topological center of certain dual algebras*, Trans. Amer. Math. Soc. **348** (1996), 1191-1212.
15. S. Mohamadzadiah, H. R. E. Vishki, *Arens regularity of module actions and the second adjoint of a derivation*, Bulletin of the Australian Mathematical Society **77** (2008), 465-476.
16. M. Neufang, *Solution to a conjecture by Hofmeier-Wittstock*, Journal of Functional Analysis. **217** (2004), 171-180.
17. M. Neufang, *On a conjecture by Ghahramani-Lau and related problem concerning topological center*, Journal of Functional Analysis. **224** (2005), 217-229.
18. J. S. Pym, *The convolution of functionals on spaces of bounded functions*, Proc. London Math Soc. **15** (1965), 84-104.
19. A. Ülger, *Arens regularity sometimes implies the RNP*, Pacific Journal of Math. **143** (1990), 377-399.
20. A. Ülger, *Some stability properties of Arens regular bilinear operators*, Proc. Amer. Math. Soc. (1991) **34**, 443-454.
21. A. Ülger, *Arens regularity of weakly sequentially compact Banach algebras*, Proc. Amer. Math. Soc. **127** (11) (1999), 3221-3227.
22. A. Ülger, *Arens regularity of the algebra $A \hat{\otimes} B$* , Trans. Amer. Math. Soc. **305** (2) (1988) 623-639.
23. P. K. Wong, *The second conjugate algebras of Banach algebras*, J. Math. Sci. **17** (1) (1994), 15-18.
24. N. J. Young *The irregularity of multiplication in group algebra* Quart. J. Math. Soc., **24** (2) (1973), 59-62.
25. Y. Zhing, *Weak amenability of module extentions of Banach algebras* Trans. Amer. Math. Soc. **354** (10) (2002), 4131-4151.

Department of Mathematics, University of Mohghegh Ardabili, Ardabil, Iran
Email address: haghnejad@aut.ac.ir